

Scope of the ATP-ubiquitin system for intracellular protein degradation

J. Saus, J. Timoneda, J. Hernández-Yago and S. Grisolia

Instituto de Investigaciones Citológicas de la Caja de Ahorros de Valencia, Amadeo de Saboya, 4, Valencia-10, Spain

Received 8 April 1982

Proteolysis

Ubiquitin

Reticulocytes

Proteases

Mitochondria

Cytosol

1. INTRODUCTION

Intracellular proteolysis and mechanisms of regulating such processes are not clear. An ATP-dependent proteolytic system has been described in rabbit reticulocytes. This system is composed of a heat-stable polypeptide of 8500–9000 M_r , designated ATP-dependent proteolysis factor I (APF-I), identified as ubiquitin [1] and detected in rat liver [2]. The substrates tested with U-ATP-protease (ubiquitin-ATP-dependent protease system) have been [^3H]globin, ^{125}I -lactalbumin, ^{125}I -lactoglobulin and ^{125}I -BSA. It is not clear if only denatured proteins are substrates from this system [3]. Therefore, we have tested whether the U-ATP-protease from rabbit reticulocytes can degrade enzymes of mitochondrial and of cytosolic location. There was no ubiquitin-ATP-dependent proteolysis with fresh ornithine transcarbamoylase (OTC), carbamoyl phosphate synthetase (CPS), glutamate dehydrogenase (GDH), pyruvate kinase (PK), lactate dehydrogenase (LDH), glucose-6-phosphate dehydrogenase (G6PDH), as assessed by activity measurements or with ^{125}I -iodinated GDH, CPS and G6PDH. Some small ATP-dependent proteolysis was observed with ^{125}I -PK. The ATP-ubiquitin system appears to be of limited scope.

2. MATERIALS AND METHODS

2.1. Enzymes

CPS from rat liver was purified and assayed for activity as in [4] and GDH from rat liver as in [5]. OTC (a gift from A. Navarro) was purified from rat liver and assayed as in [6]. GDH from beef liver, PK from rabbit muscle and G6PDH from yeast were purchased from Boehringer.

2.2. Protein iodination

Proteins were iodinated for 5 min as in [7] with Na^{125}I (13–17 mCi/ μg ; 100 mCi/ml, 250 $\mu\text{Ci}/\text{mg}$ protein). Free $^{125}\text{I}^-$ was removed by filtration using Sephadex G-25 fine equilibrated with 0.05 M sodium phosphate buffer (pH 7.5) pre-saturated with BSA. Protein fractions from the column, containing <5% of trichloroacetic acid-soluble $^{125}\text{I}^-$ were pooled (spec. act. 10^4 – 10^5 cpm/ μg protein).

2.3. Isolation and assay of components of the ubiquitin-ATP-system

Partially purified fraction I and fraction II were obtained from rabbit reticulocytes [8]. The U-ATP-protease was assayed according to [9]. Unless indicated otherwise, incubation mixtures contained in 100 μl 100 mM Tris-HCl (pH 7.6); 0.2 μmol dithiothreitol, 1 μmol phosphocreatine, 10 μg phosphocreatine kinase, 0.5 μmol MgCl_2 , 25 μg partially purified APF-I, 1.3 mg fraction II, 50 nmol ATP, and 2 μg enzyme or BSA. All incubations were for 2 h at 37°C. Loss of enzyme activity, or liberation of trichloroacetic acid-soluble radioactivity for ^{125}I -proteins was measured. Results are expressed as the percentage of trichloroacetic acid-soluble radioactivity after 2 h incubation, corrected for the percentage at zero time.

3. RESULTS AND DISCUSSION

Here, the U-ATP-protease under the conditions in [9] liberated ~ 13% of the trichloroacetic acid-soluble radioactivity from ^{125}I -BSA, in good agreement with [9]. However, when we used native enzymes, there was no or at best slight loss of activity from the enzymes tested (table 1,2). Neither

Table 1

Effect of the U-ATP-protease on several cytosolic enzymes (% proteolysis or activity after 2 h at 37°C)

Protein	- ATP	+ ATP
^{125}I -BSA (Control)	1.5	12.5
Pyruvate kinase	108	103
Lactate dehydrogenase	103	112
Glucose-6-phosphate dehydrogenase	105	105

Incubations were done as in section 2 using 10 μg enzymes listed or 2 μg ^{125}I -BSA

Table 2

Effect of the U-ATP-protease on several mitochondrial enzymes (% proteolysis or activity after 2 h at 37°C)

Protein	- ATP	+ ATP
^{125}I -BSA	1.5	12.5
Glutamate dehydrogenase (beef liver)	100 (108)	99 (91)
Carbamoyl phosphate synthetase	115 (80)	143 (83)
Ornithine transcarbamoylase	95 (123)	111 (110)

Incubations were done as in section 2 using 100 μg of the proteins listed, except for OTC where 2.5 μg were added. The figures between parenthesis were obtained using 3 μg of the enzymes listed

APF-I nor fraction II alone had any effect on the activities of the enzymes tested.

Given that only denatured and/or modified proteins can be substrates for the U-ATP-protease, ^{125}I -labelled proteins were tested. With ^{125}I -GDH there were no significant differences in the effects produced with either the complete system or the components separately, which could be ascribed to the U-ATP-protease (fig.1). However, fraction II had some proteolytic activity 'per se'. Also, when incubating ^{125}I -BSA or other labelled proteins for 2 h (without the U-ATP-protease) there was some




Fig.1. Effect of different components of the U-ATP-dependent proteolytic system on ^{125}I -BSA and ^{125}I -GDH from rat liver. Assayed as in section 2 except where indicated.

liberation (~ 2%) of trichloroacetic acid-soluble radioactivity.

When non-specific degradation due to fraction II was subtracted (see fig.2), no significant proteolysis was seen with the different enzymes tested. It should be noted that greater activities with ^{125}I -BSA as substrate have been obtained with




Fig.2. Proteolysis of BSA and of some iodinated enzymes. Assayed as in section 2 using the indicated iodinated proteins. Where indicated ATP or APF-I were excluded. Liberation of radioactivity produced by F-II alone has been subtracted from the indicated values. GDH* means glutamate dehydrogenase from beef liver.

some preparations of fraction II (23%), in agreement with [10]; however, the results with native and iodinated enzymes were also negative (not shown).

In [11] we presented evidence for the existence of both lysosomal and non-lysosomal mechanisms for mitochondrial protein degradation. The U-ATP-protease may be involved in mitochondrial protein degradation during the reticulocyte maturation process [12].

We have tested both native and partially modified (by iodination) mitochondrial enzymes as substrates for the U-ATP-protease and found no significant inactivation and/or degradation of native or iodinated mitochondrial enzymes tested with U-ATP-protease under the same conditions that cause extensive proteolysis of BSA. We also tested a number of cytosolic enzymes and again no evidence was found for inactivation with the U-ATP-protease. Thus, under these conditions, the enzymes tested are not substrates for the U-ATP-protease. The activity of U-ATP-protease is very low; i.e., ~ 0.2–0.4 µg BSA are hydrolyzed in 2 h with ~ 1.3–2 mg fraction II protein. Thus, the U-ATP-protease may be a highly specialized system and of a limited scope.

ACKNOWLEDGEMENTS

This research has been supported in part by the 'Comisión Asesora de Investigación Científica y Técnica' and the 'Fondo de Investigaciones Sanitarias' of Spain. J.S. is an 'S. Ochoa' Fellow of the Ayuntamiento de Valencia.

REFERENCES

- [1] Wilkinson, K.D., Urban, M.K. and Hass, A.L. (1980) *J. Biol. Chem.* 255, 7529–7532.
- [2] Ciechanover, A., Elias, S., Heller, H., Ferber, S. and Herskho, A. (1980) *J. Biol. Chem.* 255, 7525–7528.
- [3] Ciechanover, A., Heller, H., Elias, S., Haas, A. and Herskho, A. (1980) *Proc. Natl. Acad. Sci. USA* 77, 1365–1368.
- [4] Soler, J., Timoneda, J., DeArriaga, D. and Grisolia, S. (1981) *Biochem. Biophys. Res. Commun.* 97, 100–106.
- [5] Timoneda, J., Wallace, R. and Grisolia, S. (1981) *Biochem. Biophys. Res. Commun.* 101, 555–562.
- [6] Lusty, C., Jilka, R. and Nietsch, E. (1979) *J. Biol. Chem.* 254, 10030–10035.
- [7] McConahey, P.J. and Dixon, F.J. (1966) *Int. Arch. Allergy* 29, 185–189.
- [8] Ciechanover, A., Hod, Y. and Herskho, A. (1978) *Biochem. Biophys. Res. Commun.* 81, 1100–1105.
- [9] Herskho, A., Ciechanover, A. and Rose, J.A. (1979) *Proc. Natl. Acad. Sci. USA* 76, 3107–3110.
- [10] Wilkinson, K.D. and Audhya, T.K. (1981) *J. Biol. Chem.* 256, 9235–9241.
- [11] Grisolia, S., Knecht, E., Hernández-Yago, J. and Wallace, R. (1980) in: *Protein degradation in Health and Disease*, CIBA Found. Symp. 75, 167–188, Elsevier Biomedical, Amsterdam, New York.
- [12] Muller, M., Dubiel, W., Rathmann, J. and Rapoport, S. (1980) *Eur. J. Biochem.* 109, 405–410.